

EKHUFT MBRRACE 2020 Data

Updated October 2022







Definitions (MBRRACE)

- Stillbirth death in the uterus (unborn) after 24 weeks gestation.
- Stillbirth rate expressed per 1,000 total births
- Neonatal death baby born alive after 24 weeks gestation, but died within 28 days.
- Neonatal death rate expressed per 1,000 live births.
- Extended perinatal deaths all stillbirths over 24 weeks and all neonatal deaths, baby born alive after 24 weeks gestation, but died before 28 days of age.

MBRRACE


- Mothers and Babies Reducing Risk through Audit and Confidential Enquiries
- Based in Oxford
- Commenced 2013
- Produce yearly reports on perinatal and maternal deaths
- Subdivide and benchmark using comparator groups EKHUFT are in group "Trust with Level 3 Neonatal Unit".
- Crude data undergo "stabilising and adjustment"
- All perinatal losses from 23 weeks subject to Perinatal Mortality Review Tool – LMNS oversight
- All term neonatal deaths and intrapartum stillbirths investigated by HSIB

Adjustment Criteria

- Mother's age
- Socio-economic deprivation based on the mother's residence
- Baby's ethnicity
- Baby's sex
- Whether they are from a multiple birth
- Gestational age at birth (neonatal deaths only)

MBRRACE Comparator Group (Trusts with a level 3 NICU) – Stillbirth Rate





More than 5% higher than the group average
Rate suppressed due to small numbers

Within 5% of the group average


Between 5 and 15% lower than the group average

More than 15% lower than the group average

MBRRACE Stillbirth Rates – Crude and Adjusted


	EKHUFT Local Data			MBRRACE Data		
Year	Total Births	Total Stillbirths	EKHUFT Crude Rate per 1000	MBRRACE Crude Rate for EKHUFT	MBRRACE Rate Adjusted for EKHUFT	MBRRACE Average for comparator group
2013	7,039	24	3.41	3.58	4.28	4.75
2014	7,000	31	4.43	4.85	5.01	4.98
2015	7,063	22	3.11	3.66	4.31	4.41
2016	6,987	27	3.86	3.70	4.12	4.11
2017	6,977	21	3.01	2.72	3.82	3.95
2018	6,569	27	4.11	3.80	4.00	3.95
2019	6,413	27	4.21	4.2	4.07	3.97
2020	6,125	20	3.27	3.6	3.77	3.71
2021	6,213	25	4.02			





MBRRACE Comparator Group (Trusts with a level 3 NICU) – Neonatal Death Rate






MBRRACE Neonatal Death Rates – Crude and Adjusted


	EKHUFT Local Data			MBRRACE Data			
Year	Live Births	Total Neonatal Deaths <28 days	Crude Rate	MBRRACE Crude Rate for EKHUFT	MBRRACE Rate Adjusted for EKHUFT	MBRRACE Average for comparator group	
2013	7,015	10	1.43	1.29	1.95	2.09	
2014	6,969	14	2.01	1.86	1.93	1.97	
2015	7,041	14	1.99	1.62	2.01	2.04	
2016	6,959	20	2.87	2.57	2.53	2.10	
2017	6,956	21	3.02	3.01	2.84	2.09	
2018	6,542	11	1.68	1.68	2.08	1.92	
2019	6,386	19	2.98	2.97	2.99	1.88	
2020	6,105	7	1.15	0.99	1.53	1.67	
2021	6,188	9	1.45				





MBRRACE Comparator Group (Trusts with a level 3 NICU) – Extended Perinatal Death Rate






MBRRACE Extended Perinatal Death Rates – Crude and Adjusted



	EK	HUFT Local Da	ta	MBRRACE Data		
Year	Total Births	Total Extended Perinatal Deaths	EKHUFT Crude Rate per 1000	MBRRACE Crude Rate for EKHUFT	MBRRACE Rate Adjusted for EKHUFT	MBRRACE Average for comparator group
2013	7,039	34	4.83	4.87	5.90	6.88
2014	7,000	45	6.43	6.70	7.02	6.94
2015	7,063	36	5.10	5.27	6.30	6.44
2016	6,987	47	6.73	6.26	6.63	6.20
2017	6,977	42	6.02	5.72	6.64	6.03
2018	6,569	38	5.78	5.47	6.06	5.86
2019	6,413	46	7.17	7.16	6.91	5.78
2020	6,124	27	4.41	4.58	5.29	5.37
2021	6,212	34	5.47			



Each Baby Counts (RCOG)

- Launched by Royal College of Obstetricians & Gynaecologists. Closed in 2019
- Ambition to reduce stillbirths, neonatal deaths and brain injuries occurring during term labour
- Eligible babies: Term (37+0 weeks gestation) babies who died during labour (intrapartum stillbirth), who were born alive but died within 7 days of birth (early neonatal death) or who meet the criteria for babies born with severe brain injury (grade III hypoxic-ischaemic encephalopathy (HIE) or therapeutically cooled (active cooling only) or decreased central tone and comatose and had seizures)
- EKHUFT had the following reportable babies in each year:
 - **–** 2015: 18
 - 2016: 29
 - 2017: 19
 - 2018: 24
 - 2019: 19

Note: some babies fell into more than one category – i.e. babies with HIE who also died within 7 days